



All Databases

PubMed

Nucleotide

Protein

Genome

Structure

OMIM

PMC

Journals

Books

Search PubMed

for

Go

Clear

Limits Preview/Index History Clipboard Details

Display Abstract

Show 20

Sort by

Send to

About Entrez

Text Version

All: 1 Review: 0

Entrez PubMed

Overview

Help | FAQ

Tutorial

New/Noteworthy

E-Utilities

PubMed Services

Journals Database

MeSH Database

Single Citation Matcher

Batch Citation Matcher

Clinical Queries

Special Queries

LinkOut

My NCBI (Cubby)

Related Resources

Order Documents

NLM Catalog

NLM Gateway

TOXNET

Consumer Health

Clinical Alerts

ClinicalTrials.gov

PubMed Central

1: Anal Biochem. 2003 Nov 15;322(2):225-32.

Related Articles, Links

ELSEVIER
FULL-TEXT ARTICLE**Cellodextrin preparation by mixed-acid hydrolysis and chromatographic separation.****Zhang YH, Lynd LR.**

Thayer School of Engineering, Dartmouth College, Hanover, NH 03755, USA.

A procedure for preparation of purified cellodextrins in gram quantities was developed for use in biochemical and microbiological studies. Cellodextrins were prepared by hydrolyzing microcrystalline cellulose (Avicel) over a period of 4 to 5.5h in the presence of a mixture of 80% (v/v) concentrated hydrochloric acid (approximately 37 wt.%) and 20% (v/v) concentrated sulfuric acid (approximately 98 wt.%) at room temperature (22 degrees C). Acetone precipitation, washing ion exchange, and neutralization with barium hydroxide were used to generate a solution of mixed cellodextrins substantially free of acids and salts. Yields following hydrolysis and precipitation were approximately 0.05, approximately 0.07, approximately 0.06, and approximately 0.02 g/g cellulose for cellotriose (G(3)), cellotetraose (G(4)), cellopentoase (G(5)), and cellohexose (G(6)), respectively. Cellodextrins with degrees of polymerization from 3 to 11 were separated chromatographically using a 29 x 5-cm I.D. Bio-Rad AG50W-X4 column arranged in series with a 91 x 5-cm I.D. Bio-Gel P4 column. This two-column system was used to obtain cellodextrin preparations at 240 mg/day for G(3), 330 mg/day for G(4), 260 mg/day for G(5), and 130 mg/day for G(6), with purity >99% for G(3), G(4), and G(5) and >95% for G(6). The overall procedure achieves yields comparable to the highest previously reported, employs a separation system that can readily be reused for multiple runs, and avoids use of fuming HCl.

PMID: 14596831 [PubMed - indexed for MEDLINE]

Display Abstract

Show 20

Sort by

Send to

[Write to the Help Desk](#)[NCBI](#) | [NLM](#) | [NIH](#)[Department of Health & Human Services](#)[Privacy Statement](#) | [Freedom of Information Act](#) | [Disclaimer](#)